
Subject: Re: back-propagation with GEANE

Posted by [Lia Lavezzi](#) on Mon, 28 Jun 2010 10:35:55 GMT

[View Forum Message](#) <> [Reply to Message](#)

Hi Anastasia,

EPSIL is the boundary crossing precision in cm.

I simulated 100 events of 1 GeV/c muons at $\theta = [-10, 10]$ deg, ϕ random, with magnetic field in the usual cave (air) on a silicon plane orthogonal to the z axis, placed at 300 cm and 0.02 cm thick --> the surface should be at $300 - 0.02 = 299.98$ cm.

In the plot I attach I draw the z coordinate registered on this plane in different cases:

- 1) with the ϵ as it is on svn, 0.001 both for air and for silicon
- 2) with the $\epsilon = 0.0001$ for silicon and $= 0.001$ for air
- 3) with the $\epsilon = 0.0001$ both for air and for silicon

In the third case, where I ask for a precision of 1 μm , the registered z coordinate after simulation is closer to the real surface (by printing the values from root with the Scan function I get in the three cases: 1) 299.98025, 2) 299.98025, 3) 299.98004 cm).

Then the ϵ value has an effect, but maybe it is not enough in your case, because we are in a very ideal case and we are talking of rounding errors that are below the limit set by ϵ . Anyway it should happen that in real life the rounding errors are covered by physical effect errors and finite resolutions and you can neglect them, I hope...

Cheers,

Lia.

File Attachments

- 1) [epsilon.ps](#), downloaded 411 times
-