
Subject: Re: very high reconstructed momenta of muons
Posted by [Malgorzata Gumberidze](#) on Mon, 16 Apr 2012 22:42:33 GMT
[View Forum Message](#) <> [Reply to Message](#)

Dear All,

I have made few more test after discussions with Stefano in order to check where the high momenta tail is coming from: tracking code or Kalman (as it was suggested by Gianluigi).

What i have done is to switch on native tracking by using in the pid level:

```
PndPidCorrelator* corr = new PndPidCorrelator();  
corr->SetInputBranch("SttMvdGemTrack");  
corr->SetInputIDBranch("SttMvdGemTrackID");  
corr->SetDebugMode(kFALSE);  
fRun->AddTask(corr);
```


One can see that the tail at high momenta (up to 15GeV/c) is present also with native tracking. So it means it is not coming from genfit.

It would be nice that experts can cross check this problem in the code as well.


gosia

File Attachments


1) [muon_mom_noGen.gif](#), downloaded 871 times




2) [muon_fitStat_p_noGen.gif](#), downloaded 882 times




3) [muon_theta_p_noGen.gif](#), downloaded 848 times




4) [muon_phi_p_noGen.gif](#), downloaded 775 times



5) [phi_theta_muon_noGen.gif](#), downloaded 869 times



6) [mvdHits_p_noGen.gif](#), downloaded 862 times



7) [sttHit_p_muon_noGen.gif](#), downloaded 869 times

