
Subject: [SOLVED] Beam Smear and Cross Sections
Posted by [Michael Kunkel](#) on Tue, 24 Jul 2012 19:20:27 GMT
[View Forum Message](#) <> [Reply to Message](#)

Hello again,

Another question about beam smearing.

When I apply a beam smear of a reaction, does pluto preserve the production cross section for that specific beam energy?

I have investigated the output of the code below, and it seems the answer is no. So I ask the forum in case I am coding incorrectly.

```
double ebeam_min = 1.1725;  
double ebeam_max = 5.44575;  
PBeamSmearing *beam_smear = new PBeamSmearing("beam_smear", "Beam smearing");
```

```
TF1* beam_smear_fn = new TF1("beam_smear_fn", "1./x", ebeam_min, ebeam_max);
```

```
beam_smear->SetReaction("g + p");  
beam_smear->SetMomentumFunction(beam_smear_fn);  
makeDistributionManager()->Add(beam_smear);
```

```
((PDalitzDecay *  
makeDistributionManager()->GetDistribution("eta_dalitz"))->SetUseQED(1);
```

```
PReaction my_reaction("_P1 = 2.2","g","p","p eta [dilepton [e+ e-] g]",creator,1,0,0,0);
```

Thanks
Michael
