
Subject: Re: Bear Smear and Cross Sections
Posted by [Michael Kunkel](#) on Sun, 02 Sep 2012 14:11:19 GMT
[View Forum Message](#) <> [Reply to Message](#)

I have tried to implement what you have written so far. I have run into some more questions. I use 64 models for the energy range 1.74 - 3.33 GeV in the c.m. frame. something I find strange is that in certain energy regions, PLUTO generates the cross section oddly. In the figure below, the PLUTO generated c.m. energy is depicted.

Notice the jump in "integrated cross section" between 1.89 and 1.91 GeV. This can be seen easier in the next figure below.

However, if I look at the differential cross sections for these energy ranges, I do not see a cause for this "jump", see figure below. (The shown plots are the interpolated plots from TGraphs)

Could you please clarify what is going on here?

Also, I was hoping you would look over my macro and see if maybe there was a mistake in my syntax that might have caused this issue. I am uploading the code.

EDITED:

Furthermore, I have plotted the published lab energy vs. total cross section and compared it to what PLUTO generates. There is a scaling factor, however this is just a constant onto one of the spectrum.

As seen below, at small energies, there is a discrepancy as seen mentioned previously.

Thanks
Michael

File Attachments

- 1) [PLUTO_generated_cm_energy_smear.jpeg](#), downloaded 556 times
 - 2) [PLUTO_GEN_180_240.jpg](#), downloaded 578 times
 - 3) [diff_Xsection_187_193.jpg](#), downloaded 576 times
 - 4) [eta_XSection.C](#), downloaded 210 times
 - 5) [total_Xsection_compare.jpg](#), downloaded 580 times
-