
Subject: Problem with PndKinVtxFitter and Ks decays
Posted by [2dd180b0](#) on Wed, 28 Nov 2012 12:01:10 GMT
[View Forum Message](#) <> [Reply to Message](#)

Dear all,

I've come across a problem while trying to reconstruct the following decay:

$D0 \rightarrow Ks \pi^+ \pi^-$, $Ks \rightarrow \pi^+ \pi^-$

The procedure I use is to first use PndKinVtxFitter to fit the $\pi^+ \pi^-$ from the Ks decay to a common vertex, and then use PndKinVtxFitter to combine the fitted Ks with another $\pi^+ \pi^-$ pair, to reconstruct the D0. The problem is that the fit to the $D0 \rightarrow Ks \pi^+ \pi^-$ always returns nan for the χ^2 .

The fitting code I use is exactly analogous to that used by Marius to reconstruct $\psi(3770) \rightarrow D^+ D^-$, $D^{+-} \rightarrow K^{+-} \pi^+ \pi^-$, and we think that the problem is due to PndKinVtxFitter not being able to properly handle the neutral Ks candidate.

I've attached the analysis code and EvtGen decay file that I used for this analysis.

Best,
Sean

File Attachments

- 1) [psi3770.dec](#), downloaded 432 times
 - 2) [run_ana_dobbs.C](#), downloaded 438 times
-