
Subject: Re: CBM cocktails

Posted by [Partha Pratim Bhaduri](#) on Mon, 01 Jul 2013 11:35:24 GMT

[View Forum Message](#) <> [Reply to Message](#)

Hello Tetyana,

Thank you so much for resuming the discussion forum. I would like to propose to have a single forum like this di-lepton one, instead of having different forums for low-mass and charmonia or separate discussion forums for electrons/muons.

I have tried to estimate relative contributions to the di-muon spectrum from J/Psi decay and DrellYan process. The figure is attached where I have plotted the differential distribution of production cross sections as a function of pair rapidity in cms frame, for 25 GeV p+p collisions. Both the cross sections are calculated using LO pQCD (with inclusion of proper K factors) meaning for cc-bar I consider light quark annihilation & gluon fusion; for DY qq-bar annihilation only. The pdf set used is LO MSTW2008. For DY, the mass region integrated is 2.9 to 4.5 GeV, a region used by NA50 to report their J/Psi to DY ratio data.

Thanks to Tetyana again for posting the link for the QWG latest report. Another interesting article I recently found at arxiv regarding quarkonium data:

"A database for quarkonium and open heavy-flavor production in hadronic collisions with HepData"; hep-ex/1304.2224

Regards,

Partha

File Attachments

1) [Jpsivsdrell.eps](#), downloaded 597 times
