

---

Subject: Fast Simulation- neutral particle  
Posted by [Lu Cao](#) on Mon, 31 Mar 2014 14:24:37 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Hi all,

I try to do fast simulation with this decay chain:  $pbp \rightarrow Ds^+ Ds^-$ ;  $Ds^- \rightarrow K^+ K^- \pi^-$ ;  $Ds^+ \rightarrow e^+ \nu \eta$ ;  $\eta \rightarrow \pi^+ \pi^- \pi^0$  (gamma gamma). From the examples in macro/scrut, I got some problems for simulating the neutral channel:

1) in simulation macro:

```
// generate electro-magnetic / hadronic split offs in the EMC? switch off when running w/o EMC
```

```
Bool_t enableSplitoff = kFALSE;
```

```
//enable the producing of parametrized neutral (hadronic) split offs  
if (enableSplitoff)
```

```
fastSim->EnableSplitoffs(BaseDir+"/fsim/splitpars.dat");
```

From the comments there, if I understand correctly, the "enableSplitoff" seems should be switched on for  $\eta \rightarrow \pi^+ \pi^- \pi^0$  (gamma gamma). I tried kTRUE, but it crashed.

2) in analysis:

```
// *** name of the only PidAlgo TClonesArray in fsim  
TString pidalg = "PidChargedProbability";
```

Does it mean no neutral algorithm in fast simulation? then, how to fill the neutral list?

Thanks in advance.

Lu

---