
Subject: Re: info on charged particle simulations into NeuLAND

Posted by [Jan Mayer](#) on Fri, 10 Feb 2017 14:17:06 GMT

[View Forum Message](#) <> [Reply to Message](#)

Hi,

welcome back.

First, the missing files:

- 'r3b_cave_vacuum.geo' is simply 'r3b_cave.geo' with 'vacuum' instead of 'Air' in line 3. This does make a significant difference (as you would expect for charged particles flying through 14m of air). I'll commit it to the repository for convenience.
- 'neuland_v2_1400cm_30dp.geo.root' needs to be generated, as there are too many variations to include them all. Run

```
root -l -q ${VMCWORKDIR}/macros/r3b/geo/create_neuland_demo_geo.C(30, 1400, "v2_1400cm_30dp")
```


The macros are not completely up to date but should be enough to start looking into the matter. Here, some primary charged particles with ~same AMeV are generated.

I'm not quite sure what you want to know about the response of NeuLAND to charged particles. Due to the huge amount of material, the total energy can be obtained for many ions/energies. One can, in principle, do the full ΔE shenanigans, see the attached example for p,d,t,He-3,He-4 with fixed ~300 AMeV and an ~100-600 AMeV range.

Best,
Jan

File Attachments

1) [deltaEE-range.png](#), downloaded 867 times



2) [deltaEE.png](#), downloaded 757 times

