

---

Subject: Problems with Fitters

Posted by [Jennifer Pütz](#) on Wed, 18 Apr 2018 07:39:04 GMT

[View Forum Message](#) <> [Reply to Message](#)

---

Dear PandaRoot User,

I have some problems with the mass constraint fit.

Proton pi-

Up to now, I used the following scheme for my reconstruction (this is a simplification of what I am doing):

```
-----  
RhoMassParticleSelector* lambdaMassSelector = new RhoMassParticleSelector("lambda0",  
fm0_lambda0, 0.3);
```

```
RhoCandList piplus, antiproton, lambdabar, xiplus;  
RhoCandList LambdaBarFinal;
```

```
lambdabar.combine(piplus, antiproton);  
lambdabar.Select(lambdabarMassSelector);  
lambdabar.SetType(3122);
```

```
loop over all lambdabar candS{
```

```
  RhoKinVtxFitter vtxfitter(lambdabar[j]);  
  vtxfitter.Fit();
```

```
  double prob_vtx = vtxfitter.GetProb();
```

```
    RhoKinFitter massFitter(lambdabarVtxFit);  
    massFitter.AddMassConstraint(fm0_lambdabar0);  
    massFitter.Fit();
```

```
  double prob_mass = massFitter.GetProb();
```

```
  if (prob_vtx>0.01 & prob_mass>0.01){  
    LambdaBarFinal.Append(lambdabarVtxFit);  
  }
```

```
combine(LambdaBarFinal, piplus)
```

```
....  
-----
```

Software:

pandaRoot trunk rev. 30202

FairSoft: my16p1

FairRoot: v.17-03

I am selecting a  $\lambda$ bar by performing a vertex fit and a mass constraint fit, but pass vertex fitted candidate to the next stage of reconstruction. This candidate has to fulfill the condition, that passed both fits with a probability of more than 1%.

Now, I changed my code to pass the candidate after the mass constraint fit to the next stage of reconstruction. This change is causing a huge loss of events.

Here are some numbers of my output (the sample contains 5000 generated events):

1) pass candidate after vertex fit (but passing both cuts) to next stage

particle type | no. of cand w/o cut (mct) | final selected (mct)

$\lambda$	2469	2236	
$\lambda$ bar	2131	1663	
XiPlus	1074	722	
XiPlus Lambda K-	277		168 (after 4C-Fit)

2) pass candidate after mass constraint fit to next stage

particle type | no. of cand w/o cut (mct) | final selected (mct)

$\lambda$	2469	2236	
$\lambda$ bar	2131	1663	
XiPlus	1075	630	
XiPlus Lambda K-	233		89 (after 4C-Fit)

I have the feeling that the quality of the fitted candidates in method 2) is worse than in 1).

I also had a look at the probability distributions of the XiPlus (see attachment)

Alessandra Lai made a cross-check with her channel. She observes a difference, too. But in her sample, the effect seems to be small.

She generated 1000 events with 1) dec17 and 2) trunk rev. 30123

The following table shows the reconstruction efficiency of the full decay (all mct) with  
a) pass the vtx fitted candidate; b) pass mass fitted candidate

1a) 2.2 %  
1b) 1.8 %

2a) 9.3 %  
2b) 8.9 %

Now my question: Is there anyone of you knowing where this comes from?


Best wishes

Jenny

File Attachments


1) [vtxprob\\_passvtxfittedcand.gif](#), downloaded 497 times

## VtxFit\_prob (pass




2) [vtxprob\\_passmassfittedcand.gif](#), downloaded 475 times

# VtxFit\_prob (pass r



3) [massfitprob\\_passvtxfittedcand.gif](#), downloaded 512 times

# MassFit\_prob (pass



4) [massfitprob\\_passmassfittedcand.gif](#), downloaded 509 times

# MassFit\_prob (pass

