Subject: Re: GEM tracking Posted by Radoslaw Karabowicz on Thu, 30 Apr 2009 09:55:32 GMT View Forum Message <> Reply to Message

Problem continues...

It seems that my yesterday enthusiastic mail was a bit preliminary. I have been looking at several tracks at it looked rather ok, however when I look systematicly into the results I have realised that the fitting is still producing not optimal results. As you may remember, I was giving to the Kalman fitter a Monte Carlo momentum increased by 100Mev/c as a starting value. What I have realized is that the fitter smears the momentum with some sigma, but the reconstructed mean momentum value of many tracks still differs from the MC momenta by a value of 100MeV/c.

I have therefore made two runs of my reconstruction, once with having the start momenta given to Kalman decreased by .5 GeV/c as compared to MC truth, and in the second run I set start momenta as MC truth plus .5 GeV/c. Plots below summarize the results:

Momenta decreased: StartMom.SetMagThetaPhi(beforeMom.Mag()-0.5, beforeMom.Theta(), beforeMom.Phi());

Momenta increased: StartMom.SetMagThetaPhi(beforeMom.Mag()+0.5, beforeMom.Theta(), beforeMom.Phi());

Please take a look at the mean value of the distribution, the reco momenta is just smeard, it doesn't come back to MC truth:((.

The next thing I am planing to do is to use the tracks with high p_t. But that will come probably the next week.

yours, radek

File Attachments
1) momDiff_m5.gif, downloaded 713 times

Page 1 of 2 ---- Generated from GSI Forum



2) momDiff_p5.gif, downloaded 731 times

