
Subject: Helix and FairTrackParH
Posted by [Stefano Spataro](#) on Wed, 22 Jul 2009 15:37:02 GMT
[View Forum Message](#) <> [Reply to Message](#)

Hello,
I would like to understand the helix parameters used in FairTrackParH, which are different from the ones present in the old-style PndTpcLheTrack.

Could somebody write, according to our "standard" definition of helix parameters, how our track is represented?
The helix parameters are: x0, y0, z0, lambda, phi, charge (even if it is q/p in FairTrackParH).
How can I know the following functions?

x = x(s)
y = y(s)
z = z(s)
px = px(s)
py = py(s)
pz = pz(s)

I have checked some tracking papers but without founding this representation. This is very important for fast correlation inside the barrel.

Thanks in advance.

Subject: Re: Helix and FairTrackParH
Posted by [Alberto Rotondi](#) on Thu, 23 Jul 2009 08:24:42 GMT
[View Forum Message](#) <> [Reply to Message](#)

Dear Stefano,

the system is called Global Cartesian in the enclosed Wittek report. We call it MARS (MAster Reference System).

There are routines MARS to SD, MARS to SC and vice versa in the GEANE interface. If you have problems to use them also Lia can help.

Best regards Alberto

File Attachments

1) [wittek.pdf](#), downloaded 6005 times

Subject: Re: Helix and FairTrackParH
Posted by [Stefano Spataro](#) on Thu, 23 Jul 2009 11:00:48 GMT
[View Forum Message](#) <> [Reply to Message](#)

Hello,
thanks for the Wittek paper. I have read it and found the formula for the momenta:

$px = p \cos(\lambda) \cos(\phi)$

$$p_y = p \cos(\lambda) \sin(\phi)$$
$$p_z = p \sin(\lambda)$$

(which is also present in your tracking report).

But this is just the "static" relation between angles and momentum coordinated in a well defined point. What I need is the coordinated p_x p_y p_z along the track, starting from the first point up to the last point (where p_z remains constant, while p_x and p_y rotate). Therefore, this formula does not help me.

In Lia's thesis, page 42 (50 in the pdf), there is the following formula:

$$x(s) = x_0 + R_h[\cos(\phi_0 + h s \cos(\lambda)/R_h) - \cos(\phi_0)]$$
$$y(s) = y_0 + R_h[\sin(\phi_0 + h s \cos(\lambda)/R_h) - \sin(\phi_0)]$$
$$z(s) = z_0 + s \sin(\lambda)$$

but unfortunately nothing for momenta.

Are those parameters (ϕ_0 λ R) the same of the ones in FairTrackParH? Is it possible to use this representation as it is with our trackbase code, or maybe x_0 y_0 z_0 are calculated in a well defined point, ϕ_0 is defined in a different way, and so on?

Thanks in advance to everybody who could answer to my questions.

Subject: Re: Helix and FairTrackParH

Posted by [Gianluigi Boca](#) on Thu, 23 Jul 2009 11:44:11 GMT

[View Forum Message](#) <> [Reply to Message](#)

hi Stefano,

I don't know if this answers your question.

Find the momenta at any given point of the trajectory is rather trivial when one knows the initial momentum and the radius and center of the helix.

It is just enough to remember that for Panda $P_x^2 + P_y^2$ is constant and that the momentum in the XY plane is always perpendicular to the radius of the helix.

Does this answer your question?

Tschus und aufwiedersehen !

Gianluigi

Subject: Re: Helix and FairTrackParH

Posted by [Stefano Spataro](#) on Thu, 23 Jul 2009 13:45:19 GMT

[View Forum Message](#) <> [Reply to Message](#)

Hi,

the only point is that in this representation I do not have the center of the circle, then I have no idea on what is the ϕ of my point, and I cannot rotate. And where is defined ϕ_0 ?

I would like some explanations on this side, or better I would like to read the equations.

Subject: Re: Helix and FairTrackParH
Posted by [Gianluigi Boca](#) on Thu, 23 Jul 2009 16:15:12 GMT
[View Forum Message](#) <> [Reply to Message](#)

ok, here is the recipy.

You need to know only the starting point of the track and the starting momentum.

Then :

- 1) find the centre of the Helix circle in the XY plane by drawing a segment perpendicular to the 2-dimensional (P_x , P_y) vector. Such a segment has direction $(-P_y, P_x)$. Find the point on such segment that is a Radius distant from the initial point of the track. That is the center of the trajectory. Here there is an ambiguity that depends on the charge of the track. In other words, you must know the charge of track in order to decide on which side the center lies.
- 2) You ϕ_0 is defined by the starting point of your track with respect to any arbitrary but chosen once for all reference frame in the XY plane.

If you want you can call me on the phone (+496159711680) for better explanations.

Gianluigi

Subject: Re: Helix and FairTrackParH
Posted by [Stefano Spataro](#) on Thu, 23 Jul 2009 16:26:45 GMT
[View Forum Message](#) <> [Reply to Message](#)

Hi,

considering that there parameters are the official ones for FairTrackParH and that somebody has implemented them, I was hoping to have exactly his definition, in order to avoid misunderstanding. Without the exact definition, I cannot evaluate the equations.

Subject: Re: Helix and FairTrackParH
Posted by [Lia Lavezzi](#) on Thu, 23 Jul 2009 17:41:38 GMT
[View Forum Message](#) <> [Reply to Message](#)

Hi Stefano,

just to clarify the meaning of the different parametrizations:

- 1) in FairTrackParH you have the 5 parameters: q/p , λ , ϕ , y_{perp} , z_{perp} . q/p is the charge over momentum, λ and ϕ are the dip and polar angles which describe the direction of the particle, y_{perp} and z_{perp} are the coordinates of the point in the SC reference frame (the one perpendicular to the momentum). Using the functions of that class you can also get the position and momentum in the master reference frame;

2) in the equations you cited from my thesis, x_0 , y_0 , z_0 define the starting point of the helix, ϕ_0 is the azimuthal angle of the starting point with respect to the helix axis, λ is the dip angle and R_h is the radius of curvature of the helix.

So, I don't think you can directly apply the equations in 2) to the track from 1) since ϕ and ϕ_0 should be different: in fact ϕ_0 is calculated with respect to the helix axis and so they would coincide only if the helix axis was coincident with the beam.

If you want to apply the $x(s)$, $y(s)$, $z(s)$ equations you should choose as starting point the one you get with `GetPosition` in `FairTrackParH` and then apply them, but you have to calculate the corresponding ϕ_0 , since it is not directly given by `GetPhi` in `FairTrackParH`.

Concerning the momentum calculation I guess that once you have the helix described in the $x(s)$, $y(s)$, $z(s)$ point you can obtain the p_x and p_y coordinates by considering the transverse momentum as tangent to the track (a circle) in that point and then projecting it on the x/y directions, can't you?

If I find the exact equations for the momentum I will post a message!

Ciao,
Lia.

Subject: Re: Helix and FairTrackParH
Posted by [Stefano Spataro](#) on Thu, 23 Jul 2009 18:04:46 GMT
[View Forum Message](#) <> [Reply to Message](#)

Lia Lavezzi wrote on Thu, 23 July 2009 19:41

1) in `FairTrackParH` you have the 5 parameters: q/p , λ , ϕ , y_{perp} , z_{perp} . q/p is the charge over momentum, λ and ϕ are the dip and polar angles which describe the direction of the particle, y_{perp} and z_{perp} are the coordinates of the point in the SC reference frame (the one perpendicular to the momentum). Using the functions of that class you can also get the position and momentum in the master reference frame;

Let me correct you, in `FairTrackParH` the parameters are:

```
/** fLm = Dip angle */  
Double_t fLm;  
/** fPhi = azimuthal angle */  
Double_t fPhi;  
/** Points coordinates in SC system */  
Double_t fX_sc, fY_sc, fZ_sc;
```

Or at least these are the data members and the parameters used in the helix constructor. There is no x_{perp} , y_{perp} and ϕ_0 , but x_{sc} , y_{sc} , z_{sc} and ϕ . Apart from λ I have no idea on what the other parameters are.

Quote:

2) in the equations you cited from my thesis, x_0 , y_0 , z_0 define the starting point of the helix,

phi0 is the azimuthal angle of the starting point with respect to the helix axis, lambda is the dip angle and Rh is the radius of curvature of the helix.

So, I don't think you can directly apply the equations in 2) to the track from 1) since phi and phi0 should be different: in fact phi0 is calculated with respect to the helix axis and so they would coincide only if the helix axis was coincident with the beam.

Well, can I consider x_sc y_sc z_sc as the starting point of the helix? In this case can phi be considered as phi0?

Quote:

If you want to apply the x(s), y(s), z(s) equations you should choose as starting point the one you get with GetPosition in FairTrackParH and then apply them, but you have to calculate the corresponding phi0, since it is not directly given by GetPhi in FairTrackParH.

...

For this reason I think it would be good to have written somewhere the definition of the FairTrackParH parameters, to avoid misunderstanding (such as phi and phi0) and in order to avoid ambiguity.

Subject: Re: Helix and FairTrackParH

Posted by [Lia Lavezzi](#) on Thu, 23 Jul 2009 20:02:44 GMT

[View Forum Message](#) <> [Reply to Message](#)

Quote:Quote:1) in FairTrackParH you have the 5 parameters: q/p, lambda, phi, yperp, zperp. q/p is the charge over momentum, lambda and phi are the dip and polar angles which describe the direction of the particle, yperp and zperp are the coordinates of the point in the SC reference frame (the one perpendicular to the momentum). Using the functions of that class you can also get the position and momentum in the master reference frame;

Let me correct you, in FairTrackParH the parameters are:

```
/** fLm = Dip angle */
Double_t fLm;
/**fPhi = azimuthal angle */
Double_t fPhi;
/** Points coordinates in SC system */
Double_t fX_sc, fY_sc, fZ_sc;
```

Ok, it's just a matter of names

If you have a look in the FairTrackParH.h, the comment at the beginning contains the 5 parameters:

q/p, lambda, phi, y_perp, z_perp

In the code the name of the parameters are:

```
fLm = lambda
fPhi = phi
fX_sc = x_perp
fY_sc = y_perp
fZ_sc = z_perp
```

Concerning the names of the coordinate, let me explain: usually the SC frame coordinates are marked as "orthogonal" x, "orthogonal" y and "orthogonal" z, just because the SC frame has its yz plane orthogonal to the track momentum.

When you describe the track you need, of the 5 parameters, 1 parameter to describe the momentum magnitude (fQp is a data member of FairTrackPar), 2 to describe the direction (lambda/phi) and 2 to describe the position (y and z... call them _perp or _sc, they are the same thing).

Quote:Well, can I consider x_sc y_sc z_sc as the starting point of the helix? In this case can phi be considered as phi0?

Actually I don't think so: phi is given with respect to the z axis, while phi0 is given with respect to an axis which passes through the center of curvature of the helix; you could consider phi = phi0 only if the helix axis was the z axis. At least this is what I understand...

Quote:For this reason I think it would be good to have written somewhere the definition of the FairTrackParH parameters, to avoid misunderstanding (such as phi and phi0) and in order to avoid ambiguity.

There is a definition of the parameters in our report on GEANE, since the parameters for both FairTrackParP and FairTrackParH are the ones used there (in GEANE, I mean) in the SD and SC frame. Maybe we can add a link in the code comment to easily get it. Would it be useful?

Ciao,
Lia.

Subject: Re: Helix and FairTrackParH

Posted by [Stefano Spataro](#) on Fri, 24 Jul 2009 11:18:58 GMT

[View Forum Message](#) <> [Reply to Message](#)

Quote:

Ok, it's just a matter of names

Ok, but you can

If you have a look in the FairTrackParH.h, the comment at the beginning contains the 5 parameters:

q/p, lambda, phi, y_perp, z_perp

In the code the name of the parameters are:

fLm = lambda

fPhi = phi

fX_sc = x_perp

fY_sc = y_perp

fZ_sc = z_perp

It is a matter of names only if you know what those names mean (not my case) And once they are also called in a different way, the amount of confusion in my mind increases.

However, it seems I do not need those parameters for my helix, even because all the definitions are to move from one system to another, but what I need is how the position/momentum varies along the track.

I have found a document in internet that explains what I need, to characterize my track:

center of the helix (to calculate ρ , and ϕ_0)

radius of the helix (to calculate ρ)

p_t (to calculate k)

λ

Of these 5 params, I have already λ , p_t , and the radius can be calculated from p . I am only missing the center of the helix, that can be calculated with the suggested method of Gianluigi.

I will try and cross my fingers hoping that it will work.