Subject: eta_c with event mixing for STT
Posted by Dima Melnychuk on Mon, 29 Aug 2011 22:35:11 GMT

View Forum Message <> Reply to Message

Dear colleagues,

I have analaised the eta_c data produced on the GRID with event mixing, and at the moment I still do not understand the results.

But I just want to share some plots to show the current status.

So multiplicity of charged tracks:

Only 8% of events have >=4 tracks which is much lower than 72% without event mixing.

Invariant mass of phi and eta_c with preselection on phi mass window:

Here there is no indication of phi or eta_c peak.

And aplying MC Truth PID does not change this distribution much.

And finaly only 4 events out of 100k pass the selection cuts.

So the results still have to be understood.

Dima

File Attachments
1) n_charged_stt.png, downloaded 1073 times

n charged

nc	
Entries	97890
Mean	1.303
RMS	1.526

2) m_nocuts_stt.png, downloaded 1131 times

φ: m(K+ K-) (no cuts)

h_mphi_nocuts	
Entries	74483
Mean	1.134
RMS	0.1372

η_c m(φ,φ), (no cuts)

h_etac_nocut	
Entries	41551
Mean	2.735
RMS	0.2301

Subject: Re: eta_c with event mixing for STT
Posted by Dima Melnychuk on Fri, 02 Sep 2011 09:37:40 GMT
View Forum Message <> Reply to Message

Dear colleagues,

Here I update plots on eta_c reconstruction with STT and event mixing with runs mix301, mix311, mix321, mix331.

There is an improvement comparing to previous results but the results are still far from reasonable.

Multiplicity of charged tracks:

23% of events have >=4 tracks in comparing to 8% in previous run and 72% without event mixing.

And invariant mass of phi and eta_c with preselection on phi mass window:

Here phi peak appears but still no indication of eta_c peak.

Finally 36 events were reconstructed from 100 k.

Dima

File Attachments

φ: m(K+ K-) (no cuts)

h_mphi_nocuts	
Entries	145772
Mean	1.164
RMS	0.1402

η_c m(φ,φ), (no cuts)

h_etac_nocut	
Entries	27904
Mean	2.77
RMS	0.2324

Subject: Re: eta_c with event mixing for STT
Posted by Dima Melnychuk on Fri, 02 Sep 2011 15:26:39 GMT
View Forum Message <> Reply to Message

Just for comparison here I present invariant mass distributions for eta_c and phi with the same version of STT software (august11).

This run is produced with cleanup code and eta_c reconstruction efficiency is 7.8%

Dima

File Attachments
1) m_nocuts_nomix.png, downloaded 798 times

φ: m(K+ K-) (no cuts)

h_mphi_nocuts	
Entries	261042
Mean	1.179
RMS	0.173

η_c m(φ,φ), (no cuts)

h_etac_nocut	
Entries	40495
Mean	2.917
RMS	0.1509

Subject: Re: eta_c with event mixing for STT
Posted by Dima Melnychuk on Thu, 08 Sep 2011 00:48:23 GMT
View Forum Message <> Reply to Message

Hi all,

With the latest run eta_c events produced with STT with background mixing are reconstructable.

Multiplicity of reconstructed tracks is rather high:

With rough pre-selection on phi mass invariant mass distributions looks like:

So there is small indication of eta_c peak.

But after the final selection:

eta_c is reconstructed with efficiency 3.3%

Dima

File Attachments
1) n_charged_stt_mix.png, downloaded 911 times

2) m_nocuts_stt_mix.png, downloaded 977 times

φ: m(K+ K-) (no cuts)

h_mphi_nocuts	
Entries	840065
Mean	1.174
RMS	0.1446

η_c m(φ,φ), (no cuts)

h_etac_nocut	
Entries	933156
Mean	2.786
RMS	0.2247

3) m_final_vtx_stt_mix.png, downloaded 936 times

φ: m(K+ K-)

h_mphi_final_vtx	
Entries	67118
Mean	1.036
RMS	0.0294

η_c m(φ,φ), (cut on φ mass)

h_etac_phimass_vtx	
Entries	15014
Mean	2.966
RMS	0.0782