Subject: Problem of energy loss of proton in FairROOT Posted by Genie on Mon, 26 Jan 2015 14:51:13 GMT View Forum Message <> Reply to Message

Hello.

I'm writing a package for a TPC experiment using FairROOT. Recently we found that the energy loss of proton and alpha doesn't match to the TRIM/LISE++ result.

So, I posted a thread in GEANT4 electromagnetic process mailing list,

but it comes to my mind that it might be more related to the FairROOT forum so I registered.

I put the copy of the post in GEANT4 electromagnetic process mailing list.

If you have any comments or recommendation or hints, please let me know. Anything would be helpful

Thank you in advance. Genie

Link: http://hypernews.slac.stanford.edu/HyperNews/geant4/get/emprocess/1333.h tml

Hello.

I'm currently writing a simulation package for an experiment.

Prior to using it, we're checking if the energy loss of proton and alpha is the same as the other simulation.

We're comparing GEANT4 energy loss data with TRIM/LISE++ data.

Attached links are proton and alpha energy loss plot with TRIM/LISE++ result.

Attachment:

http://hypernews.slac.stanford.edu/HyperNews/geant4/get/AUX/2015/01/22/1 5.07-13310-0atGMsC.png

http://hypernews.slac.stanford.edu/HyperNews/geant4/get/AUX/2015/01/22/1 5.07-54093-KDTIfHd.png

As you can see, in relatively low energy, energy loss curves doesn't match each other. We couldn't find what the problem is.

Currently, we're writing our code on top of FairROOT package. So, GEANT4 is steered over G4VMC interface. Cuts used for running simulation is following.

If you have any clue or comments or recommendation, please don't hesitate to mail us. Anything could be helpful

Thank you in advance.

Best regards, Genie

g4config.in: /mcPhysics/rangeCuts 0.001 mm /cuts/setLowEdge 10 eV /cuts/setHighEdge 1 MeV q4Config.C: TG4RunConfiguration\* runConfiguration = new TG4RunConfiguration("geomRoot", "QGSP BERT HP EMY", "stepLimiter+specialCuts+specialControls"); TGeant4 -> SetMaxNStep(1.E7); SetCuts.C gMC->SetProcess("PAIR",1); /\*\* pair production \*\*/ gMC->SetProcess("COMP",1); /\*\* Compton scattering \*\*/ gMC->SetProcess("PHOT",1); /\*\* photo electric effect \*\*/ gMC->SetProcess("PFIS",1); /\*\* photofission \*\*/ gMC->SetProcess("DRAY",1); /\*\* delta-ray \*\*/ gMC->SetProcess("ANNI",1); /\*\* annihilation \*\*/ gMC->SetProcess("BREM",1); /\*\* bremsstrahlung \*\*/ gMC->SetProcess("HADR",1); /\*\* hadronic process \*\*/ gMC->SetProcess("MUNU",1); /\*\* muon nuclear interaction \*\*/ gMC->SetProcess("DCAY",1); /\*\* decay \*\*/ gMC->SetProcess("LOSS",1); /\*\* energy loss \*\*/ gMC->SetProcess("MULS",1); /\*\* multiple scattering \*\*/ Double t cutC = 1.0E-5; //Double t cutB = 1.E4; Double\_t cutB = -1; Double t cut1 = 1.0E-8; // [GeV] Double\_t cutp = 0.01; // [GeV] Double\_t tofmax = 1.E10; // seconds gMC->SetCut("CUTGAM",cutC); /\*\* gammas (GeV)\*/ gMC->SetCut("CUTELE",cut1); /\*\* electrons (GeV)\*/ gMC->SetCut("CUTNEU",cutC); /\*\* neutral hadrons (GeV)\*/ gMC->SetCut("CUTHAD",cutC); /\*\* charged hadrons (GeV)\*/ gMC->SetCut("CUTMUO",cutC); /\*\* muons (GeV)\*/ gMC->SetCut("BCUTE",cutB); /\*\* electron bremsstrahlung (GeV)\*/ gMC->SetCut("BCUTM",cutB); /\*\* muon and hadron bremsstrahlung(GeV)\*/ gMC->SetCut("DCUTE",cutB); /\*\* delta-rays by electrons (GeV)\*/ gMC->SetCut("DCUTM",cutB); /\*\* delta-rays by muons (GeV)\*/ gMC->SetCut("PPCUTM",cutp); /\*\* direct pair production by muons (GeV)\*/ gMC->SetCut("TOFMAX",tofmax); /\*\*time of flight cut in seconds\*/

Subject: Re: Problem of energy loss of proton in FairROOT Posted by StefanoSpataro on Mon, 26 Jan 2015 18:03:43 GMT View Forum Message <> Reply to Message Subject: Re: Problem of energy loss of proton in FairROOT Posted by Genie on Tue, 27 Jan 2015 14:54:15 GMT View Forum Message <> Reply to Message

Thank you for the comment!

I tried as you recommended, but unfortunately it doesn't change anything.

Page 3 of 3 ---- Generated from GSI Forum