

---

Subject: Different x-y resolutions

Posted by [Simone Bianco](#) on Tue, 19 Jul 2011 11:16:58 GMT

[View Forum Message](#) <> [Reply to Message](#)

---

Dear Tracking experts,

I am simulating simple events with 4 pions (2pi+ and 2 pi-) with a momentum of 1GeV/c, which are propagated from (0,0,0). I am generating these particles with an homogeneous distribution in phi and in theta (where theta is restricted to the range [20,140]°).

The simulations are performed with all the detectors in, but I set as active only the MVD, the Stt, the GEMs and the forward tracer.

I perform my reconstruction using:

PndMvdRiemannTrackFinderTask,PndSttTrackFinderReal,PndSttMvdTracking,PndRecoKalmanTask,PndMCTrackAssociator,PndPidCorrelator (with pion hypothesis),PndPidIdealAssociatorTask.

At the end of this chain I am running some analysis using the PndVtxPoca vertex finder.

I fill histograms with the three components of the reconstructed vertices. As you can see in the attached figures I get a significantly different sigmas for x and y.

X always looks worse (even with different momenta and theta ranges).

I have already tried with different topologies like using two particles instead of four, or using pions of the same charge, but I always get worse results for X.

I tried with different vertexing tools (POCA, Ralf's fast fitter, another implementation of the POCA) and I always noticed this behaviour. It really seems to come from previous steps since if I try to rotate of 90° the tracks just before running the POCA vertex finder I obtain symmetrical results (worse Y than X, see the second figure attached).

Does any of you have an idea about possible reasons of this effect?

Thanks in advance.

Cheers,

Simone


---

### File Attachments


1) [Vtx-0-0-0-1GeV-4Pions.png](#), downloaded 1305 times

---

# Vertex



2) [Rotation-Vtx-0-0-0-1GeV-4Pions.png](#), downloaded 1294 times



Subject: Re: Different x-y resolutions  
Posted by [Stefano Spataro](#) on Tue, 19 Jul 2011 11:25:32 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Have you tried the PndKinVtxFitter? There I do not see strong differences.

---

Subject: Re: Different x-y resolutions  
Posted by [Simone Bianco](#) on Tue, 19 Jul 2011 16:10:04 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

I have not done it yet. I will try this now and report about the results I obtain. Thanks for your hint

---

Subject: Re: Different x-y resolutions  
Posted by [Simone Bianco](#) on Tue, 19 Jul 2011 19:52:09 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---



Hi again, this is what I get using the PndKinVtxFitter:

X is still significantly worse than Y

#### File Attachments

1) [PndKinVtx.png](#), downloaded 1114 times

---




---

Subject: Re: Different x-y resolutions

Posted by [Jens Sören Lange](#) on Wed, 20 Jul 2011 06:40:37 GMT

[View Forum Message](#) <> [Reply to Message](#)

Hi Simone,

one idea: which field are you using?

PndMultifield("FULL") or PndMultifield("SOLENOID") only?

(i.e. can you maybe try to switch off the dipole field?)

best regards, Soeren

Subject: Re: Different x-y resolutions  
Posted by [Simone Bianco](#) on Wed, 20 Jul 2011 09:13:16 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Hi Soeren,

I have never tried this before, so I am now surprized to see that setting in the simulation macro:

```
PndMultiField *fField= new PndMultiField("SOLENOID");  
fRun->SetField(fField);
```

I get really bad results: I nearly never get all the four pions reconstructed and anyway their momentum is not the one set in the particle generator. This sounds really strange to me since I am shooting the pions in the theta range [20,140], so I would expect the dipole not to play any role in their reconstruction.

What am I doing wrong?

### File Attachments

---

1) [runSimPGun.C](#), downloaded 437 times

---

---

Subject: Re: Different x-y resolutions  
Posted by [Jens Sören Lange](#) on Wed, 20 Jul 2011 14:22:25 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Hi Simone,

1.) my idea, that it could have something to do with the B field, came from the X and Y differences which were observed for luminosity monitor (which was to my knowledge the only time, that any azimuthal asymmetry was reported). Now I re-checked, and I found that I was wrong: it was not the dipole field generating the effect, but the aperture. See

[http://forum.gsi.de/index.php?t=tree&goto=9206&rid=0&S=77db77666b257eeb980ffbd17eec1d35&srch=Jost#msg\\_9206](http://forum.gsi.de/index.php?t=tree&goto=9206&rid=0&S=77db77666b257eeb980ffbd17eec1d35&srch=Jost#msg_9206)

However, same as the dipole field, the effect should be forward peaked.

2.) HOWEVER, your result shows, that there is an effect even in your polar angle range.

After all, there might be an effect which was probably never seen before because you are looking at very small track DCA resolutions (50-60 um), but the B field grid is (if I remember correctly) in the central region 4 cm point-to-point distance (with linear interpolation between the points).

I checked field/PndMultiField.cxx

```
if (Map=="FULL") {
```

```
    PndTransMap *map_t= new PndTransMap("TransMap", "R");  
    PndDipoleMap *map_d1= new PndDipoleMap("DipoleMap1", "R");
```

```

PndDipoleMap *map_d2= new PndDipoleMap("DipoleMap2", "R");
PndSolenoidMap *map_s1= new PndSolenoidMap("SolenoidMap1", "R");
PndSolenoidMap *map_s2= new PndSolenoidMap("SolenoidMap2", "R");
PndSolenoidMap *map_s3= new PndSolenoidMap("SolenoidMap3", "R");
PndSolenoidMap *map_s4= new PndSolenoidMap("SolenoidMap4", "R");

AddField(map_t);
AddField(map_d1);
AddField(map_d2);
AddField(map_s1);
AddField(map_s2);
AddField(map_s3);
AddField(map_s4);

}else if (Map="DIPOLE") {
    PndDipoleMap *map_d1= new PndDipoleMap("DipoleMap1", "R");
    PndDipoleMap *map_d2= new PndDipoleMap("DipoleMap2", "R");

    AddField(map_d1);
    AddField(map_d2);

}else if (Map=="SOLENOID") {

    PndSolenoidMap *map_s1= new PndSolenoidMap("SolenoidMap1", "R");
    PndSolenoidMap *map_s2= new PndSolenoidMap("SolenoidMap2", "R");
    PndSolenoidMap *map_s3= new PndSolenoidMap("SolenoidMap3", "R");
    PndSolenoidMap *map_s4= new PndSolenoidMap("SolenoidMap4", "R");

    AddField(map_s1);
    AddField(map_s2);
    AddField(map_s3);
    AddField(map_s4);
}

```

so when you change from "FULL" to "SOLENOID", it means:  
it is the SAME solenoid maps, but only the dipole map and the trans map IN ADDITION (!).

In the moment, I don't understand why you see such a strong effect for the dipole field. The best would be to switch off the B field completely, but then of course the track fitter does not work for straight tracks. I have the feeling that the track fitter also uses the field map, and then fails if B field in Geant and fitter are different. But I don't know where to set the field map for the fitter. I hope that someone else has a clever idea.

Soeren

---

Subject: Re: Different x-y resolutions  
 Posted by [Elisa Fioravanti](#) on Wed, 20 Jul 2011 15:12:23 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Hello Simone,


I'm studying the pi+pi-pi+pi- analysis with the dataset available on GRID (pandaroot version july11).

I would like to show you the resolution that I have with the PndKinVtxFitter, in the STT case. As you can see, in my case the X and Y resolution are almost the same.

best  
Elisa

## File Attachments

1) [vertex.png](#), downloaded 857 times



---

Subject: Re: Different x-y resolutions

Posted by [Simone Bianco](#) on Wed, 20 Jul 2011 15:17:11 GMT

[View Forum Message](#) <> [Reply to Message](#)

---

Hi Elisa,

I suppose there you have a forward boost for the system, which theta range do your pions span? Are these tracks measured with the mvd/stt/gems only or you use the forward tracker as well?

Cheers,

Simone

---

Subject: Re: Different x-y resolutions  
Posted by [Elisa Fioravanti](#) on Wed, 20 Jul 2011 15:41:16 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Hello Simone,

I used EvtGen and the four pions are simulated with phase space distribution.  
And the tracks are measured with stt/mvd/gem only

Elisa

---

---

Subject: Re: Different x-y resolutions  
Posted by [Simone Bianco](#) on Wed, 20 Jul 2011 16:34:24 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Dear Soeren,

I performed one test: I ran the simulation with the full field, but setting the fBeamMom to a small value, so that the map chosen for the dipole field is the least intense. With this setup I see no differences with the case in which fBeamMom was bigger (leading to a different dipole map). I would expect this change to show some effects if the x-y difference is in some way due to the dipole, but this doesn't seem to be the case. I will keep on investigating, thanks for your help!

Best regards,

Simone

---

---

Subject: Re: Different x-y resolutions  
Posted by [Tobias Stockmanns](#) on Thu, 21 Jul 2011 07:25:18 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Hi Simone,

it is possible to set the magnetic field to a constant value without using the field maps. This is done in PndConstField.

Maybe you can try this method to exclude any effects of an asymmetric magnetic field.

Cheers,

Tobias

---

---

Subject: Re: Different x-y resolutions  
Posted by [Simone Bianco](#) on Thu, 21 Jul 2011 15:54:46 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Thanks Tobias, the constant field worked fine. Using the same parameters as before I got really similar results:

So, this should exclude effects of asymmetries in the magnetic field.

I tried to select smaller theta ranges and what I notice is that in the "forward" part everything works fine. For example here you see what I obtained (always with four pions of 1 GeV/c) limiting the theta range to the range [30,40]°:

Increasing theta the situation gets worse. This is what I obtained in the range [60,70]:

Cheers,

Simone


PS: all these plots were obtained using the POCA vertex finder

#### File Attachments


---

1) [ConstField\\_20\\_140deg.png](#), downloaded 1151 times

# Vertex



2) [ConstField\\_30\\_40deg.png](#), downloaded 1205 times



3) [ConstField\\_60\\_70deg.png](#), downloaded 1096 times

