

---

Subject: Bremsstrahlung and neutrals mergeing  
Posted by [Klaus Götzen](#) on Tue, 06 May 2014 12:18:53 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Hi all,

I just wanted to inform you that I implemented a neutral (cluster) mergeing and bremsstrahlung losses for electrons in the Fast Sim in trunk and scrut14. As default these features are switched off and have to be enabled in the fast sim macro with

```
// enable the merging of neutrals if they have similar direction  
fastSim->MergeNeutralClusters();
```

```
// enable bremsstahlung loss for electrons  
fastSim->EnableElectronBremsstrahlung();
```

The parametrization for the cluster mergeing was motivated by Ronalds talk (see [https://panda-wiki.gsi.de/foswiki/pub/Computing/Minutes28April2014/evo14\\_0428RK.pdf](https://panda-wiki.gsi.de/foswiki/pub/Computing/Minutes28April2014/evo14_0428RK.pdf)) to roughly match the efficiency on page 11.

The following plot shows the  $J/\psi \rightarrow e^+ e^-$  mass distribution (from  $\psi' \rightarrow J/\psi \pi^+ \pi^-$  events) without (blue) and with (red) bremsstrahlung:

This plot shows the  $\pi^0$  reco efficiency depending on the opening angle (here from  $p\text{bar}p \rightarrow \pi^+ \pi^- \pi^0$  events):


You might give it a try and check, whether it behaves like expected. In case of your positive feedback we can also enable the features as default.

Best,  
Klaus


---

### File Attachments

1) [j2e\\_brems.gif](#), downloaded 1587 times



2) [pi0\\_merge.gif](#), downloaded 1374 times




---

Subject: Re: Bremsstrahlung and neutrals mergeing  
 Posted by [Elisa Fioravanti](#) on Wed, 07 May 2014 11:43:41 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Hello Klaus,

I tried the bremsstrahlung for the simulation of the decay:

chi\_cj->J/psi gamma

with J/psi->e+e-

I first selected a J/psi from e+e- candidates, and then

I did a 4C fit to the chi\_cj candidates.

If I look to the probability of the fit I obtained this distribution:

with a peak at zero.

But when I simulated the same decay mode, using the same macro,  
 but without the bremsstrahlung, after the 4C fit I obtained this  
 probability distribution:


uniform distributed from 0 to 1.

Do you have an idea on what's going on?


Thanks  
Elisa

## File Attachments

1) [bremm.png](#), downloaded 1126 times



2) [nobremm.png](#), downloaded 1045 times



---

Subject: Re: Bremsstrahlung and neutrals merging  
Posted by [Klaus Götzen](#) on Wed, 07 May 2014 11:57:12 GMT  
[View Forum Message](#) <> [Reply to Message](#)

Hi Elisa,

I think this is a realistic behaviour in the sense, that the candidates with a large bremsstrahlung

loss don't fit the 4C very well and therefore get low probability. You might check with full mc, whether you see there also a peak for MC truth matched candidates close to 0 ...

Best,  
Klaus

---

---

Subject: Re: Bremsstrahlung and neutrals mergeing  
Posted by [StefanoSpataro](#) on Wed, 07 May 2014 11:59:08 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Hi Klaus,  
in this plot it seems that half of the counts (3000) have prob 0. Maybe it is a too high loss in efficiency.

---

---

Subject: Re: Bremsstrahlung and neutrals mergeing  
Posted by [Klaus Götzen](#) on Wed, 07 May 2014 12:22:47 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Hi,

if you take a look to full sim, it seems quite comparable (at least the shape of the tail). But I could put a parameter controlling at least the probability that bremsstrahlung takes place. This for now is constant about 30%, which I estimated from full sim single particles.

Best,  
Klaus

---

---

Subject: Re: Bremsstrahlung and neutrals mergeing  
Posted by [Ralf Kliemt](#) on Wed, 07 May 2014 13:30:35 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

StefanoSpataro wrote on Wed, 07 May 2014 13:59Hi Klaus,  
in this plot it seems that half of the counts (3000) have prob 0. Maybe it is a too high loss in efficiency.

Since there are two "electrons" involved in the decay, we have a probability for at least one Bremsstrahlung photon of  $(1-0.7^2)=0.51$ . The technically interesting question is: Is it real hard zeros or just very small values (as i would expect).

Cheers  
Ralf

---

Subject: Re: Bremsstrahlung and neutrals mergeing  
Posted by [Elisa Fioravanti](#) on Wed, 07 May 2014 13:46:23 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Hello,

I tried to do the same plot in order to see how many events are exactly zero and the result is 1448 events. I attached the plot.

Elisa

### File Attachments

---

1) [zero.png](#), downloaded 1348 times

