
Subject: [FIXED] Problem with mass constraint fit for two gamma

Posted by [donghee](#) on Mon, 30 Jun 2014 08:56:57 GMT

[View Forum Message](#) <> [Reply to Message](#)

Hi

I found a strange behaviour with mass constrain fit.

For charged combination (here is e.g. $D0 \rightarrow K\pi^+$), chi2 distribution of mass constrain fit are similar and seems to be OK for both full and fast simulation.

An issue about mass constraint fit has been fixed in few weeks ago. (see message #16518 : Problems with PndKinFitter::AddMassConstraint).

However one more problem is still remaining.

For neutral combination (here is e.g. $D0 \rightarrow \gamma\gamma$), chi2 distribution and also probability distribution of mass constrain fit are too different in the case of full and fast simulation at scrut14. I have correctly delta function for invariant mass for gamma gamma combination after mass constraint fit.

Only problem is the two low chi2 distribution(distributed in below 0.1) in the full simulation.

full_gg_mass_fit_results_to_forum_charged.eps show the chi2 distribution of the mass fit for $D0 \rightarrow K\pi^+$ in two cases(fast and full).

full_gg_mass_fit_results_to_forum_neutral.eps show the chi2 distribution of the mass fit for $D0 \rightarrow \gamma\gamma$.

Below is the accessor what I have commonly used for full and fast with scrut14 version.

```
//Pnd Kin Mass Fitter
PndKinFitter mfitter(dp[j]);      // instantiate the kin fitter in
mfitter.AddMassConstraint(D0Mass); // set 4 constraint
mfitter.Fit();                    // do fit
double chi2_m = mfitter.GetChi2(); // get chi2 of fit
double prob_m = mfitter.GetProb(); // access probability of fit
RhoCandidate *dpfit_mass = dp[j]->GetFit(); // get fitted D0
```

I don't understand why the chi2 distributions are so different for the neutral combination in both case(full and fast)?

I assume that some factors are missing or NDF is handled wrongly for the neutral combination case in the full simulation .

Best wishes,
Donghee

File Attachments

- 1) [full_gg_mass_fit_results_to_forum_charged.eps](#), downloaded 516 times
- 2) [full_gg_mass_fit_results_to_forum_neutral.eps](#), downloaded

Subject: Re: Problem with mass constraint fit for two gamma

Posted by [Lu Cao](#) on Mon, 30 Jun 2014 14:51:42 GMT

[View Forum Message](#) <> [Reply to Message](#)

Hi Donghee and all,

this situation is the same in reconstructing $\pi^0 \rightarrow \gamma \gamma$, as I reported before.

In the fast sim, the invariant mass spectrum of the combination of two photons has a "nice" distribution peaked at π^0 mass, so it's understandable that the mass constraint fit can work as expected. In the full sim, the invariant mass distribution is filled with a significant "background" which may cause problems of convergence in mass constraint fit, e.g. too small χ^2 , prob squeezed to 1.


Since the `PndKinFitter::AddMassConstraint()` works fine for charged particles as well as the neutral in the fast sim, I'd like to guess the tool itself has no problem but to understand why the two-photon invariant mass distribution has so much "background" in the full sim. Is there any possible incorrect combinatorics, or where're the extra photons coming from?

Donghee, could you also show the $\gamma\gamma$ invariant mass spectrum in your case?

Best regards,
Lu

File Attachments

1) [mass_two-photon.png](#), downloaded 555 times



Subject: Re: Problem with mass constraint fit for two gamma
Posted by [StefanoSpataro](#) on Mon, 30 Jun 2014 15:51:59 GMT
[View Forum Message](#) <> [Reply to Message](#)

I repeat for the nth time: the EMC covariance matrix in full simulation was never filled properly, and there is currently nobody from EMC groups working to have a proper implementation. In fast everything is fine since the error matrix is what you impose in the simulation.

Lu, your plot is wrong, if you use a mass constraint fit then you force your background to stay inside your peak. The fit is not done to remove background, but to select your events once you have already removed the background.

Subject: Re: Problem with mass constraint fit for two gamma
Posted by [donghee](#) on Tue, 01 Jul 2014 08:12:52 GMT
[View Forum Message](#) <> [Reply to Message](#)

Hi Stefano,

I will be a volunteer to help this issue. (will try, but cannot guarantee about completeness)
First of all, I have to know what is the difference for the covariance matrix in EMC part, and better to compare it between fast and full.
Where is the EMC covariance matrix in the full simulation?

Best wishes,
Donghee

Subject: Re: Problem with mass constraint fit for two gamma
Posted by [StefanoSpataro](#) on Tue, 01 Jul 2014 08:24:29 GMT
[View Forum Message](#) <> [Reply to Message](#)

In input folder.

Subject: Re: Problem with mass constraint fit for two gamma
Posted by [donghee](#) on Tue, 01 Jul 2014 08:35:58 GMT
[View Forum Message](#) <> [Reply to Message](#)

Hi Stefano,

In input folder, there are couple of emc_matrix as a root format, but I cannot read it.
Who produce them?
Is this file some kind of mapping according to momentum, theta, and phi?

Best wishes,
Donghee

Subject: Re: Problem with mass constraint fit for two gamma
Posted by [StefanoSpataro](#) on Tue, 01 Jul 2014 08:37:31 GMT
[View Forum Message](#) <> [Reply to Message](#)

Please contact Dima Melnychuk.

Subject: Re: Problem with mass constraint fit for two gamma
Posted by [donghee](#) on Tue, 01 Jul 2014 08:57:29 GMT
[View Forum Message](#) <> [Reply to Message](#)

Hi Lu,

Left is for the D0 (gamma gamma) and right is after mass constraint fit.


As you mentioned, chi2 is squeezed to 0 also in my case.
But I guess that the background issue is not connected with this problem of mass fitter.
We need to solve matrix problem of EMC in the case of full simulation, then chi2 distribution

should be reasonable.

Best wishes,
donghee

File Attachments

1) [cl.png](#), downloaded 849 times



Subject: Re: Problem with mass constraint fit for two gamma
Posted by [Stefano Spataro](#) on Tue, 01 Jul 2014 08:59:06 GMT
[View Forum Message](#) <> [Reply to Message](#)

Can you show the chi2 distributions?

Subject: Re: Problem with mass constraint fit for two gamma
Posted by [Dima Melnychuk](#) on Tue, 01 Jul 2014 09:06:49 GMT
[View Forum Message](#) <> [Reply to Message](#)

Hi Donghee,

The EMC error matrix is accessed via class `/emc/EmcReco/PndEmcErrorMatrix.cxx` and it is used in `PndPidCorrelator` to actually set error matrix for neutral candidates. The parameters used in this parametrization are stored in `/input/emc_error_matrix_default.root` and

/input/emc_error_matrix_1.root

Error matrix is parametrized on energy and position of the cluster with independent parametrizations for Barrel Forward and Backward endcaps. I have attached the presentation from Mark Pelizaeus (I hope he does not mind) who did it in old framework and I adapted it for pandaroot. I remember he had another presentation on the subject but I cannot find it now.

emc_error_matrix_default.root contain parameters obtained by Marc in old framework and in emc_error_matrix_1.root there are parameters which I calculated for one emc geometry ("emc_module12.dat", "emc_module3new.root", "emc_module4_StraightGeo24.4.root", "emc_module5_fsc.root"), but it is not exactly the one used by default now. For forward endcap it was modified to "emc_module3_2012_new.root".

There are several macros related to emc error matrix in /macro/emc/dedicated/

error_matrix_data_production.C - produces photons in the whole emc range to calculate error matrices

error_matrix_fit.C - to actually calculate error matrices, fit them, and store parameters in PndEmcErrorMatrixParObject containers.

fill_error_matrix_param.C - to fill PndEmcErrorMatrixParObject container with given parameters.

I just found that in PndEmcErrorMatrix.cxx the emc_error_matrix_default.root was used and not emc_error_matrix_1.root for default geometry and I will correct it now (within one hour) and you can try if this can solve the problem.

Otherwise it makes sense to start with some QA macros to test pull distribution of pi0 before and after 4C-fit as suggested in Mark's presentation and if there is actually a problem with error matrices, to produce photons, re-calculate error matrices, parametrize them and test again.

Dima

File Attachments

1) [M.Pelizaeus, emccov_4cfit_pbm.pdf](#), downloaded 376 times

Subject: Re: Problem with mass constraint fit for two gamma

Posted by [donghee](#) on Tue, 01 Jul 2014 09:08:46 GMT

[View Forum Message](#) <> [Reply to Message](#)

Hi Stefano,

Here is the chi2 distribution for D0->gamma gamma mass constraint fit.

(detail plots are in the file at first posting of this title)


Black histogram is the all candidates and dashed blue is for the MC truth matched one. I assume that the chi2 values are normalized some factor.

I will discuss with Dima.

Best wishes,
Donghee

File Attachments

1) [c2.png](#), downloaded 887 times



Subject: Re: Problem with mass constraint fit for two gamma
Posted by [Dima Melnychuk](#) on Tue, 01 Jul 2014 09:58:26 GMT
[View Forum Message](#) <> [Reply to Message](#)

I just updated the PndEmcErrorMatrix.cxx and now the emc_error_matrix_1.root error matrix parametrization is used for default emc geometry instead of emc_error_matrix_default.root. There is a chance that it can improve the results.

Subject: Re: Problem with mass constraint fit for two gamma
Posted by [donghee](#) on Tue, 01 Jul 2014 12:17:18 GMT
[View Forum Message](#) <> [Reply to Message](#)

Hi Dima,

Thank you for your kind explanation.
I think that you have already made all relevant study for error matrices.

It means that in fast simulation the error matrix "emc_error_matrix_1.root" has been accessed correctly but in full simulation old parameterization emc_error_matrix_default.root was used up to now.

Did you make parametrizations for Barrel, Forward endcap, Backward endcap, forward EMC, all 4 parts separately?

what is the difference between "emc_module3new.root" and "emc_module3_2012_new.root". Your parametrization is based on the former, but the default mapper in PANDAroot (version=1) is the latter.

Do we need a re-calculation of error matrix using modul "emc_module3_2012_new.root"?

If you think that we need some huge photon production in each kinematic binning and more study for error matrices, please let me know I am ready to do that.

Best wishes,
Donghee

Subject: Re: Problem with mass constraint fit for two gamma

Posted by [donghee](#) on Wed, 02 Jul 2014 09:57:56 GMT

[View Forum Message](#) <> [Reply to Message](#)

Hi Dima,

I try again with modified PndEmcErrorMatrix.cxx and the emc_error_matrix_1.root, but chi2 distribution is not improved.

It is still squeezed as before. I am wondering why doesn't change anything?

Either We need to recalculation for error matrice or have to find some bugs during the filling the error of matrices in the code.

Best wishes,
Donghee

Subject: Re: Problem with mass constraint fit for two gamma

Posted by [StefanoSpataro](#) on Wed, 02 Jul 2014 10:01:52 GMT

[View Forum Message](#) <> [Reply to Message](#)

Hi Donghee,

I try to repeat again: the current EMC error matrices were not calculated with the current EMC geometry and reconstruction code, they should be recalculated.

Subject: Re: Problem with mass constraint fit for two gamma
Posted by [Dima Melnychuk](#) on Wed, 02 Jul 2014 10:26:50 GMT
[View Forum Message](#) <> [Reply to Message](#)

Hi,

It's pity that it does not help, i.e. usage of `emc_error_matrix_1.root` instead of `emc_error_matrix_default.root`.

Comment to Stefano, that `emc_error_matrix_1.root` was calculated for geometry which is different from current only for forward endcap and modification in reconstruction code should not be (?) relevant for this case.

Comment to Donghee

> It means that in fast simulation the error matrix "`emc_error_matrix_1.root`" has been accessed correctly but in full simulation old parameterization `emc_error_matrix_default.root` was used up to now.

No, in fast simulation the error matrices were calculated/parametrized independently.

> Did you make parametrizations for Barrel, Forward endcap, Backward endcap, forward EMC, all 4 parts separately?

There is a separate parametrization for each EMC part but I calculated them in one macro.

But the conclusion is that error matrices should be recalculated at least for modified forward endcap, but parametrization/fit should be checked anyway, since I do not suppose that only forward endcap introduces problems.

So it makes sense to start with some QA macro to look for pull distribution for π^0 before/after 4C-fit and only later to recalculate error matrices.

Dima

Subject: Re: Problem with mass constraint fit for two gamma
Posted by [donghee](#) on Wed, 02 Jul 2014 10:28:02 GMT
[View Forum Message](#) <> [Reply to Message](#)

Hi Stefano,

Dima suggested simply solution as like this, but it doesn't make sense...it is pity.
I will discuss with him and try to produce correct matrix with current geometry.

Best wishes,

Subject: Re: Problem with mass constraint fit for two gamma

Posted by [StefanoSpataro](#) on Wed, 02 Jul 2014 11:24:27 GMT

[View Forum Message](#) <> [Reply to Message](#)

Hi Dima,

If I remember correctly, last year there was the open question about non-uniformity light collection and electronic noise, and I believe the error matrices were calculated before we were settings the proper flags. In this sense I believe that this has screwed a bit the situation.

Subject: Re: Problem with mass constraint fit for two gamma

Posted by [donghee](#) on Wed, 02 Jul 2014 22:29:39 GMT

[View Forum Message](#) <> [Reply to Message](#)

Hi Dima,

I am starting to look the re-etsimation of the error matrices for EMC.

Clearly We need some modifications of QA macros.

And I found an unclear thing for the photon production, why all other detectors are turned off when the photons are produced?

In order to take into account material effect, MVD,STT,Drc and so on every detector system have to be put into the simulation.

Best wishes,

Donghee

Subject: Re: Problem with mass constraint fit for two gamma

Posted by [Dima Melnychuk](#) on Thu, 03 Jul 2014 11:29:49 GMT

[View Forum Message](#) <> [Reply to Message](#)

Hi Donghee,

It's actually a question I do not have an answer, to include all the other subdetectors or not when calculationg error matrices.

Other subdetectors can cause a preshower with some probablility and photons reconstructed after preshower will have worse resolution, even if they are reconstructed at all.

And I do not know if to include them in error matrix calculation or not.

But I do not have strong arguments against it, so you can use the whole PANDA for error matrix calculation if you want.

Dima

Subject: Re: Problem with mass constraint fit for two gamma

Posted by [donghee](#) on Thu, 03 Jul 2014 21:57:10 GMT

Hi Dima,

I have some technical question for ths simulation.

We need to know the energy of emc cluster in this task.

There are two quantity, one is the raw energy and the other one is corrected energy by `GetEnergyCorrected()` from clustering.

In my understanding, corrected energy is the energy deposit in the cluster after calibration.

We are using normally corrected energy in the analysis level, so I assume that we have to use corrected one for the error estimation too.

Otherwise, error matrix should be overestimated without correction(calibration).

Best wishes,
Donghee

Subject: Re: Problem with mass constraint fit for two gamma
Posted by [Dima Melnychuk](#) on Fri, 04 Jul 2014 09:23:08 GMT
[View Forum Message](#) <> [Reply to Message](#)

Hi Donghee,

Actually I do not see a problem using raw energy for error matrix assuming the same energy is used for parametrization and calculation of error matrices. I.e. error matrix is calculated for the proper raw energy (in `PndPidCorrelator`) but may be the error itself will be expressed in non-corrected energy units but I think it's a second order effect here.

On the other hand using corrected energy will complicate the code. The method `GetEnergyCorrected()` of the `PndEmcCluster` is obsolete and in principle should be removed. You obtain corrected energy with `PndEmcClusterCalibrator` class, as for example (`fEmcCalibrator->Energy(cluster)`) and using corrected energy will create dependence of `PndEmcErrorMatrix` class on `PndEmcClusterCalibrator`, which is not a problem if it's necessary, but I am not sure it is.

Dima

Subject: Re: Problem with mass constraint fit for two gamma
Posted by [Stefano Spataro](#) on Sun, 06 Jul 2014 04:09:50 GMT
[View Forum Message](#) <> [Reply to Message](#)

Before starting to calculate matrices, if I were in you I would check first the actual code.

First, is the information stored in the neutral candidate correct? Or is maybe the error matrix somehow bugged? Maybe something in the initialization went wrong? When I launch a pid macro i can see the following error:

-I- FairRunTimeDB::InitContainer() PndEmcErrorMatrixPar

[ERROR] init() PndEmcErrorMatrixPar not initialized

I could suspect the par is not properly loaded.

Then, if I launch the photon simulation I can see fCov[15] is not set (it is at zero) while fErrP7[28] has strange numbers, also negative (-30?). If I do GetErrorP7() I obtain something at 10^6 !

Are we sure the correct data members are properly filled? Which variables are needed really for the kinematic fitters? Maybe we are just filling the wrong thing, or maybe we are filling it in the wrong way (i.e. position inverted with momentum, mass at the beginning and not at the end, the usual stuff).

Could you please someone do such checks?

Subject: Re: Problem with mass constraint fit for two gamma
Posted by [donghee](#) on Mon, 07 Jul 2014 09:04:57 GMT

[View Forum Message](#) <> [Reply to Message](#)

Hi Stefano and Dima,

I am now looking your posting for the neutral candidate in pid correlator.
Launching the PID macro, Error message doesn't have serious meaning because we didn't store the emc parameter (PndEmcErrorMatrixPar) into the data base.

Quote:

```
// Get Emc error matrix parameter container
fEmcErrorMatrixPar = (PndEmcErrorMatrixPar*) db->getContainer("PndEmcErrorMatrixPar");
```

We can ignore this alarm, or we can make inactive thie matrix parameter container before initialization, because we will access error matrix from the input reading file later on.

Quote:

```
void PndPidCorrelator::SetParContainers() {
...
// Get Emc error matrix parameter container
fEmcErrorMatrixPar = (PndEmcErrorMatrixPar*) db->getContainer("PndEmcErrorMatrixPar");
...
}
```

I find a potential problem for miss matching of neutral candidate.

When we construct a neutral candidate using EMC, we build candidates from "bump" information based, not EMC "cluster".

But error calculation is previously made by EMC cluster.

I don't know exactly which one has more fundamental information and is placed more high rank.

Best wishes,
Donghee

Subject: Re: Problem with mass constraint fit for two gamma
Posted by [Dima Melnychuk](#) on Mon, 07 Jul 2014 09:13:31 GMT
[View Forum Message](#) <> [Reply to Message](#)

Hi Donghee,

there is no problem using cluster for error calculation and applying them to bumps since PndEmcBump is in principle a PndEmcCluser except for the cases when PndEmcCluser is produced from pi0 and therefore cluster has two local energy maxima and is divided into two bumps, otherwise EmcBump is a copy of EmcCluster.

Dima

Subject: Re: Problem with mass constraint fit for two gamma
Posted by [donghee](#) on Mon, 07 Jul 2014 11:54:37 GMT
[View Forum Message](#) <> [Reply to Message](#)

Hi Dima and Stefano,

Error matrix in the contructuion of nueutral candidate has a form with 7X7 matrices in total 3 categories.

I have seen/learn follow setting for neutral cand. with PndPidCandidate.
position error set to be 0.
position-momentum covariance set to be also 0.
momentum(px,py,pz,E) error has been evaluated by module and cluster.

I am wondering the term in position-momentum covariance must be 0 for neutral case? or I am now misunderstanding for this concept.

Best wishes,
Donghee

Subject: Re: Problem with mass constraint fit for two gamma
Posted by [StefanoSpataro](#) on Mon, 07 Jul 2014 12:03:56 GMT
[View Forum Message](#) <> [Reply to Message](#)

In theory the kinematic fitter should not use position neither pos errors.

Subject: Re: Problem with mass constraint fit for two gamma
Posted by [StefanoSpataro](#) on Wed, 16 Jul 2014 15:49:46 GMT
[View Forum Message](#) <> [Reply to Message](#)

Dear Donghee,
Could you please update the Emc folder of the trunk and try again? I think I found and fixed the bug.

Subject: Re: Problem with mass constraint fit for two gamma
Posted by [donghee](#) on Wed, 16 Jul 2014 18:28:56 GMT
[View Forum Message](#) <> [Reply to Message](#)

Dear Stefano,

Thanks a lot.
I submit a job with this new EMC reconstruction part.

Probably, I focused on the wrong direction before getting your e-mail.
You can see what I was suspicious about and discussion with Dima at today morning.

See you soon,

Hi Dima,

I am produing emc data(photon) to get error matrices according current geometries.
Everythings is more or less smooth.

I found that the position resolution of backward endcap are really bad, if I compare with forward end cap.
Did you see also same behaviour?

I assume that our main problem is the scaleing factor in the PndEmcErrorMatrix.cxx
We are calculating the position error at z or R=100cm when we estimate errors and save it from the analysing photon data.
Then in PndEmcErrorMatrix we make a rescale with $54/100 \times \text{error}$, which 54 is the radius of barrel for example and 100 is previously given in the error matrix data that have to be subtracted during the access this information at analysis (fit) level.

But I think that this is something wrong. we don't need any scale factor. All error are estimated by differences at certain position.
It has been already taken into account during transformation in which is TMatrix trans(4,3).

I am trying to do with following way.
In PndEmcErrorMatrix, I change
from
`pos1Err *=scaleFactor`
`pos2Err *=scaleFactor`

to
pos1Err *=1/100
pos2Err *=1/100

let see what should happen.
Donghee

Subject: Re: Problem with mass constraint fit for two gamma
Posted by [StefanoSpataro](#) on Wed, 16 Jul 2014 20:56:58 GMT
[View Forum Message](#) <> [Reply to Message](#)

Hi,
the mine was a bug, but the matrices need to be recalculated anyway. I hope that with my correction the chi2 will assume something closer to what we should expect, but for the correct analysis we need better matrices for sure.
No idea about the scaling factor you are speaking about. I can try to take a look tomorrow.


Subject: Re: Problem with mass constraint fit for two gamma
Posted by [donghee](#) on Thu, 17 Jul 2014 08:29:07 GMT
[View Forum Message](#) <> [Reply to Message](#)

Dear Stefano and Dima,


I am very happy to inform you that the problem is finally solved.
Test two plot show the chi2 distribution for D0(gammagamma) mass constrain fit before and after fixed error matrix in PndEmcErrorMatrix.
One need to re-run pid correlator to have correct error matrix during the analysis.
In addition, I will update soon more precise estimation of error matrices according current EMC geometry & will report about it in our PANDArOOT meeting.
Thanks again to Stefano for your valuable effort and great help.

File Attachments

1) [chi2_for_D0_before.png](#), downloaded 726 times



2) [chi2_for_D0_after.png](#), downloaded 838 times



Subject: Re: Problem with mass constraint fit for two gamma
Posted by [StefanoSpataro](#) on Thu, 17 Jul 2014 08:34:21 GMT
[View Forum Message](#) <> [Reply to Message](#)

Very good, I will copy this modification also to the scrut14.
Do you want to report then next wednesday?

Subject: Re: Problem with mass constraint fit for two gamma
Posted by [donghee](#) on Thu, 17 Jul 2014 08:59:10 GMT
[View Forum Message](#) <> [Reply to Message](#)

Dear Stefano,

I am pleasure to do that.
I will present about update of EMC error matrix with new geometry including some diagnostics(mass constraint fit) from pi0 and D0 data".

Best wishes,
Donghee

Subject: Re: Problem with mass constraint fit for two gamma
Posted by [Lu Cao](#) on Thu, 17 Jul 2014 11:20:46 GMT
[View Forum Message](#) <> [Reply to Message](#)

Hi Stefano,

why my plot is wrong?? It's the invariant mass distribution of two-photon BEFORE any fittings.
Where's the problem in this plot?
Sorry for delayed response.

Best,
Lu

Subject: Re: Problem with mass constraint fit for two gamma
Posted by [StefanoSpataro](#) on Thu, 17 Jul 2014 11:34:46 GMT
[View Forum Message](#) <> [Reply to Message](#)

It seemed like an old plot with mass after and before the fit. If this is for two different samples, then maybe it is ok. I don't remember which was the discussion exactly 20 days ago.
I would suggest to close this topic and to open some new one, since here there are two many ongoing discussions.
